Mr David Gordon MB ChB, MRCS, MD, FRCS (Tr & Orth)

Consultant Orthopaedic Surgeon

www.davidgordonortho.com





Spire Bushey Hospital

Private Practice Personal Assistant: Helen Sellars T/F: 01582-622-219 E: helen@davidgordonortho.com

NHS Secretary: Jacqueline Andrews T: 01582-497-066

Pain Diary Following Injection

Date _			Joint :	L / R	Ankl	le /	M	TPJ / PI	antar Fa	scia /
Medication administered				Hydrocortistab /			Kenalogue / Ma			arcain
Please circling the appropriate number representing the severity of your pain at each time point.										
Before Injection										
0	1	2	3	4	5	6	7	8	9	10
No Pa	in									Worst pain ever
Immediately after injection (within 15 minutes)										
0	1	2	3	4	5	6	7	8	9	10
No Pa	in									Worst pain ever
2 Hours after injection										
0	1	2	3	4	5	6	7	8	9	10
No Pa	in									Worst pain ever
End of the day of injection										
0	1	2	3	4	5	6	7	8	9	10
No Pa	in									Worst pain ever
End of the 2 weeks after injection										
0	1	2	3	4	5	6	7	8	9	10
No Pain									Worst pain ever	

Advice after the Injection

You can put as much weight as is comfortable through the foot. Remove the plaster after about 2 hours. You may experience some increase in pain within 24 hours, this will settle. The injection may take up to 7 days to work.

Please bring this form with you to your next clinic appointment